

****TITLE****

*ASP Conference Series, Vol. **VOLUME**, **PUBLICATION YEAR***

****EDITORS****

The evolution of radio galaxy emission line regions: Breaking the P-z degeneracy

K. J. Inskip, M. S. Longair

Astrophysics Group, Cavendish Laboratory, Cambridge, CB3 0HE

P. N. Best

Institute for Astronomy, Edinburgh, EH9 3HJ

Abstract. The ionization and kinematic properties of the emission line regions of three subsamples of 6C and 3CR radio galaxies have been compared. The degeneracy between redshift and radio power is broken, and the relative importance of radio power (P), radio size (D) and redshift (z) for the emission line region properties is determined.

1. Introduction

The emission line regions (ELRs) of high redshift radio galaxies are generally more extensive than those of lower redshift sources. Spectroscopic studies of 3CR sources at $z \sim 1$ (Best, Röttgering & Longair 2000a, 2000b) reveal that the size of the radio source plays an important role in determining the dominant ionization mechanism. The emission lines of small sources are best explained by the predictions of shock ionization associated with the passage of the radio cocoon. Larger sources usually appear photoionized by the central AGN, and typically exist in a higher ionization state than their smaller counterparts.

Studies of 3CR radio galaxies reveal considerable changes with redshift, with the lower redshift/radio power sources (degenerate in the flux limited 3CR sample) generally having much less extreme kinematics than sources at $z \sim 1$ (Baum & McCarthy 2000; Best et al 2000b). This degeneracy can be broken by means of a comparison with other radio galaxy surveys, e.g. the 6C sample.

2. Breaking the redshift–radio power degeneracy

We have studied a complete subsample of 8 6C galaxies within the redshift range $0.85 < z < 1.3$ (Inskip et al, *in prep*). These are a factor of ~ 6 lower in radio power than 3CR sources at the same redshift. The ionization and kinematic properties of the ELRs of these galaxies have been compared with those of the $z \sim 1$ 3CR subsample of Best et al (2000a), and also with lower redshift 3CR sources matched in radio power to the 6C $z \sim 1$ subsample. Low redshift ELR kinematic data were obtained from Baum & McCarthy (2000); ionization data were obtained from Tadhunter et al (1998). This combination of samples allows us to break the P–z degeneracy effectively.




Figure 1. These plots display the variation in maximum velocity FWHM with (a - left) D and (b - right) P for three radio galaxy subsamples. 3CR sources at $z \sim 1$ are denoted by crosses, and 6C sources at $z \sim 1$ by filled triangles. The stars denote the low redshift 3CR data. (Filled stars represent the low redshift 3CR sources with comparable radio powers to the 6C subsample.)

Fig. 1 displays kinematic data for three subsamples, plotted against D and P. Principal component analysis and partial rank correlation tests on these data sets have been used to determine the relative importance of P, D and z on the changing ionization and kinematic properties of radio galaxy ELRs. For larger radio galaxies, the gas kinematics become less extreme, and the ionization state increases, as has been found in previous studies (e.g. Best et al 2000b). In addition to this, we also find that ionization state is strongly dependent on P but *not* z , with the more powerful radio sources existing in higher ionization states than less powerful sources independent of redshift. On the other hand, the FWHM of the ELRs is correlated most strongly with redshift, as well as with radio power. The most extreme kinematics are observed in powerful radio sources at high redshift.

From this analysis we conclude that:

1. AGN properties vary little with redshift, and the power/age of a radio source is the determining factor in the ionization mechanism of the emission line gas.
2. The gas kinematics points to significant evolution of the host galaxy and ELR properties with redshift, in addition to their dependency on the properties of the radio source.

References

- Baum S.A., McCarthy P.J., 2000, ApJ, 119, 2634
 Best P.N., Röttgering H.J.A., Longair M.S., 2000a, MNRAS, 311, 1
 Best P.N., Röttgering H.J.A., Longair M.S., 2000b, MNRAS, 311, 23
 Inskip K.J., Best P.N., Rawlings S., Cotter G., Longair M.S., Röttgering H.J.A., Eales S.A., in prep.
 Tadhunter C.N., Morganti R., Robinson A., Dickson R., Villar-Martín M., 1998, MNRAS, 298, 1035